

Original Research

# The Dorsiflexion Range of Motion Screen: A Validation Study

Phillip J Plisky<sup>1</sup>, Garrett S Bullock<sup>2</sup>, Mary Beth Garner<sup>1</sup>, Risa Ricard<sup>1</sup>, Josh Hayden<sup>1</sup>, Bethany Huebner<sup>1</sup>, Kate Schwartzkopf-Phifer<sup>1</sup>, Kyle Kiesel<sup>1</sup>

<sup>1</sup> Department of Physical Therapy, University of Evansville, <sup>2</sup> Nuffield Department of Orthopaedics, Rheumatology, and Musculoskeletal Sciences, University of Oxford

Keywords: ankle, dorsiflexion, movement system, screen

<https://doi.org/10.26603/001c.21253>

---

## International Journal of Sports Physical Therapy

Vol. 16, Issue 2, 2021

---

### Background

Limited ankle dorsiflexion (DF) is associated with ankle sprains and other lower extremity injuries. Current ankle measurements can be laborious to perform in an athletic environment.

### Purpose

The purpose of this study was to determine the reliability and discriminant validity of a novel closed-chain ankle DF ROM test, the standing ankle dorsiflexion screen (SADS).

### Study Design

Reliability and validity study

### Methods

Thirty-seven healthy subjects participated in the study. Two raters measured closed-chain ankle DF range of motion (ROM) using a modified lunge position with an electronic inclinometer. Four raters measured ankle DF using the SADS. Reliability was calculated using intraclass correlation coefficients (ICC) and kappa coefficients for the raters using an electronic inclinometer and the SADS scale, respectively. An independent t-test compared the SADS categories of “behind” and “beyond” to the modified lunge test ROM ( $p < 0.05$ ).

### Results

Excellent ICC values (0.95 [95% CI (0.92,0.97)]) and high kappa values were observed (0.61-0.81), with high percent agreement (86-94%). There was a significant difference in ankle DF ROM between the nominally scored “behind” and “beyond” categories, regardless of rater or trial analyzed (behind:  $41.3^\circ \pm 4.7^\circ$ ; beyond:  $51.8^\circ \pm 6.1^\circ$ ,  $p < 0.001$ ).

### Conclusions

The SADS was observed to have excellent interrater reliability and high discriminant validity. Furthermore, there was a distinct closed chain ankle DF ROM difference between the “behind” and “beyond” SADS nominal scores.

### Clinical Relevance

The SADS can be used as a quick and efficient closed chain ankle DF ROM screen.

### Level of Evidence

2b

## INTRODUCTION

Athletic lower extremity injuries are common,<sup>1-3</sup> with ankle sprains being the most prevalent.<sup>3,4</sup> Nearly 75% of athletic ankle sprains are recurrent,<sup>5</sup> burdening the sports medicine provider and health care system.<sup>6,7</sup> Research has suggested that limited ankle dorsiflexion (DF) range of motion (ROM) increases ankle and overall lower extremity injury risk.<sup>8-11</sup> In addition, limited ankle DF is a common injury sequela. Therefore, sports medicine providers have a need for easy to use on-field ankle screens.<sup>9,12</sup>

Ankle DF ROM has conventionally been assessed in the open-chain position; however, open-chain ankle testing has poor reliability.<sup>13</sup> Furthermore, athletic movement and competition are performed in the closed-chain position.<sup>13,14</sup> Normal ankle DF allows for lower extremity advancement, running, and proper jump landing.<sup>11,15</sup> Thus, performing closed-chain ankle measurements allows for more functional clinical testing.<sup>10,16</sup> Previous closed-chain ankle DF testing has been performed in the half kneeling position<sup>17</sup> and in a modified lunge position both with high reliability and validity.<sup>18,19</sup> However, these tests require other devices, such as an inclinometer, which are not always readily available.<sup>20</sup> This decreases the utility of those tests, as well as the ability of sports medicine providers to effectively and efficiently screen ankle DF. A screen is used to quickly identify if there is a potential problem whereas a measure requires equipment and gives a numerical result. As a result, there is a need for an ankle DF screen that requires minimal equipment and can be implemented quickly and efficiently. Researchers have examined the reliability and validity of a novel ankle DF screen, but used half kneeling dorsiflexion as the referent standard.<sup>21</sup> Since the ankle dorsiflexion screen is in the standing lunge position, additional analysis comparing measure in that position is warranted.

The purpose of this study was to determine the reliability and discriminant validity of a novel closed-chain DF ROM test, the standing ankle dorsiflexion screen (SADS). It was hypothesized that the SADS would have high reliability and discriminant validity.

## METHODS

### SUBJECTS

A convenience sample of university students was utilized. Subjects were recruited using fliers on a university campus. To be included in the study, subjects needed to be over 18 years of age and ambulatory without an assistive device. Exclusion criteria consisted of participants with a previous lower extremity orthopedic surgery, current pain or injury, or diagnosed neurological disease. Informed consent was obtained from each subject prior to data collection. The University of Evansville's Institutional Review Board approved study procedures.

### MODIFIED LUNGE ROM MEASUREMENT

The modified lunge position (Figure 1) was used for the discriminant measurement.<sup>18,19</sup> Closed-chain ankle DF ROM



**Figure 1: Standing modified lunge ankle DF ROM testing position with inclinometer 2cm below tibial tuberosity.**

was measured using an electronic inclinometer (Clinometer Android App version 2.4 by Plaincode™ on Samsung Galaxy s9) in a standing modified lunge position with the subjects in a tandem heel to toe stance (Figure 1).<sup>18,19</sup> For balance, participants held a dowel rod in the contralateral hand. The inclinometer was placed two centimeters below the inferior aspect of the tibial tuberosity on the back lower-limb. Prior to ankle DF testing, subjects were instructed to drop their back knee as far down and forward as possible while taking the back knee as far as possible beyond the back toes, without lifting the back heel. The raters then recorded the dorsiflexion measurement in degrees from tibial vertical for each trial.<sup>17</sup>

### STANDING ANKLE DORSIFLEXION SCREEN

The SADS was completed using the same position for the modified lunge measurement. The back ankle DF was scored by identifying how far forward the back knee moved in relation to the front medial malleolus. The SADS was scored on an ordinal scale of three categories, *behind*, *within*, and *beyond* (Figure 2).

### ANKLE RANGE OF MOTION MEASUREMENTS AND SCREENING

Seven raters were utilized for all measurements; each rater was a physical therapist who specialized in outpatient orthopedics and sports medicine. All raters were trained prior to ankle DF measurements that consisted of demonstration and verbal instruction by an instructor who helped develop the screen and had 20 years of orthopedic physical therapy experience. Following the initial demonstration, each rater

**Table 1: Modified Lunge Test Reliability**

	MWLT Kappa CI <sub>95</sub> †	MWLT % Agreement
Rater 1 to Rater 2	.69 (.51-.78)	87
Rater 1 to Rater 3	.71 (.54-.88)	88
Rater 1 to Rater 4	.67 (.49-.85)	87
Rater 2 to Rater 3	.61 (.40-.82)	86
Rater 2 to Rater 4	.70 (.51-.89)	90
Rater 3 to Rater 4	.81 (.65-.97)	94

MWLT = Modified Weighted Lunged Test  
 †CI<sub>95</sub> = 95% Confidence Interval


performed at least three trials with feedback from the instructor. This was repeated and compared to the instructors' measurements until the instructor determined that the measurement was being taken according to the instructions.

Upon arrival of each subject, their age, height, and weight were recorded. Each subject was randomly allocated through a coin flip to first be measured with the modified lunge or SADS. The modified lunge ROM and SADS category measurements were taken two times per ankle with five minutes between measurements to prevent a treatment effect.<sup>22</sup> Two raters measured ankle DF ROM using the modified lunge position. Four raters measured ankle DF via the SADS categories of *behind*, *within*, and *beyond*. All raters were blinded to other rater's measurements.

**STATISTICAL ANALYSES**

Descriptive statistics (age, height, weight) were calculated for subjects using means ± standard deviations. Each ankle was analyzed independently. The intraclass correlation coefficient (ICC 2,2) with a 95% confidence interval was used to determine interrater reliability for goniometric ROM measurements where a score of less than 0.40 was deemed poor, 0.40 to 0.59 moderate, 0.60 to 0.74 good, and greater than 0.75 excellent.<sup>23</sup> Kappa coefficients were utilized to determine interrater reliability for the SADS categories, where <0.40 was deemed poor, 0.40-0.59 moderate, 0.60-0.74 good, and >0.75 excellent.<sup>23</sup>

The modified lunge measurements in degrees of dorsiflexion ROM were then averaged between trials. The SADS *behind* and *within* categories were dichotomized into one category (*behind*), while *beyond* remained the same. Discriminant validity of the SADS categories was assessed with an independent t-test (p<0.05), using the mean values on the modified lunge ROM of the *behind* and *beyond* categories. A significant difference in modified lunge ROM between the *behind* and *beyond* categories would indicate that the screen can discriminate between individuals with satisfactory ankle dorsiflexion and individuals with limited ankle dorsiflexion. All analyses were conducted in SPSS 24 (IBM SPSS Statistics for Windows, Version 24.0. Armonk, NY: IBM Corp).



**Figure 2: Standing Ankle Dorsiflexion Screen testing position. Color indicating patella position at maximal ROM. Green = beyond malleolus. Yellow = within malleolus. Red = behind malleolus**

**RESULTS**

The convenience sample consisted of 37 ambulatory subjects (74 ankles), including 27 females (age: 23 ± 1.1 years; height: 167.2 ± 10.2 cm; body mass: 66.9 ± 11.5 kg) and 12 males (age: 22.9 ± 0.7 years; height: 178.3 ± 9.1 cm; mass: 77.6 ± 10.3 kg). The interrater reliability was excellent for the modified lunge electronic inclinometer measurements, with an ICC value of 0.95 (95% CI: 0.92-0.97). The kappa coefficient was high for the SADS categories (range: 0.61-0.81), with percent agreement ranging from 86 to 94% (Table 1).

There was a statistically significant difference in closed-chain ankle DF ROM between the *behind* and *beyond* cate-

gories regardless of rater or trial analyzed (behind:  $41.3^\circ \pm 4.7^\circ$ ; beyond:  $51.8^\circ \pm 6.1^\circ$ , SEM 1.36,  $p < 0.001$ ).

## DISCUSSION

Limited ankle DF is an injury risk factor and common sequela after injury.<sup>3,4</sup> Many ankle DF measurement techniques are not performed in a standing position and require expensive equipment.<sup>13,20</sup> Thus, there is a need for an easy to administer ankle screen that requires no equipment. The purpose of this study was to determine the reliability and discriminant validity of a novel closed-chain DF ROM screen, the SADS. Supporting the hypothesis, the SADS was observed to have excellent interrater reliability and discriminant validity. Furthermore, there was a distinct DF ROM difference between the subjects with behind and beyond SADS nominal scores.

The SADS demonstrates discriminant validity. In this study, a digital inclinometer was used. Venturni et al.<sup>24</sup> compared the reliability of ankle DF measurement using both a standard bubble goniometer and a digital inclinometer. The results indicated high (ICC=0.83) interrater reliability for the digital inclinometer and moderate (ICC=0.72) interrater reliability for the bubble goniometer.<sup>24</sup> This corroborates the decision to study the SADS discriminant validity with an inclinometer. A significant difference in modified lunge ROM between the subjects in the *behind* and *beyond* categories was observed, though the mean for the *behind* category was higher than expected.

Using a functional closed-chain position for ankle screening was found to have high reliability compared to the modified lunge tests. This supports previous research in which a functional closed-chain position was found to have the greatest reliability.<sup>22,25,26</sup> Munteanu et al.<sup>22</sup> found that ankle DF measurements in a knee extended weight-bearing position had high interrater and intrarater reliability in both novice and experienced raters.<sup>22</sup> Five positions for measuring DF were compared by Krause et al.,<sup>25</sup> with the modified lunge having the greatest intrarater (ICC=0.88-0.89) and interrater (ICC=0.82) reliabilities. The authors concluded that the modified lunge position may best assess end-range ankle DF.<sup>25</sup> This highlights the fact that performing an ankle screen in a functional closed-chain position may best assess potential differences in ankle DF ROM.

Closed-chain ankle DF measurements in the modified lunge were similar to previous studies.<sup>8,21,27,28</sup> The SADS *behind* score mean was 41 degrees. Malliaras et al.<sup>8</sup> found an increased risk of patellar tendinopathy was associated with less than 45 degrees of closed-chain DF in volleyball players. Athletes with repeated ankle sprains have been observed to have decreased ankle DF ROM.<sup>14,29</sup> The SADS *beyond* score mean was 51.8 degrees. Driller et al.<sup>27</sup> ob-

served that healthy subjects demonstrated over 50 degrees of ankle DF during the weight-bearing lunge test. Dill et al.<sup>28</sup> found that healthy subjects with normal ankle motion had 51 degrees DF in the weight-bearing lunge position, while limited ankle DF subjects had 39 degrees of ankle DF. In the study previously examining the reliability and validity of the SADS using half kneeling dorsiflexion as the referent standard, there were differences between DF ROM measurements.<sup>21</sup> The half kneeling DF ROM measurements were  $33.5 \pm 2.0$  degrees for behind,  $38.6 \pm 1.2$  degrees for within, and  $43.0 \pm 0.78$  degrees for beyond compared to the standing lunge measurement of behind  $41.3^\circ \pm 4.7^\circ$  and beyond  $51.8^\circ \pm 6.1^\circ$ .<sup>21</sup> However, the reliability was similar in this study ranging from 0.61 to 0.81 with percent agreement from 86% to 96%.<sup>21</sup>

There were limitations in this study. This study was limited to subjects who were all injury-free at time of testing and were college aged students. However, these participants may have had chronic ankle instability, chronic Achilles tendinopathy, or plantar fasciitis, which may demonstrate long-term ankle or foot impairment. Therefore, the generalization of results outside these ages and to subjects current ankle and/or lower extremity injuries is not possible. Further research should include diverse age ranges, sports, and individuals with ankle and/or lower extremity injury to increase external validity. Additionally, prospective studies are required to determine the SADS injury risk identification ability.

## CONCLUSION

The SADS is a reliable and valid ankle screen for assessing closed-chain ankle DF ROM. Ankle DF ROM differences between the subjects with SADS nominal scores *behind* and *beyond*, were significantly different. The SADS can be used as a quick and efficient closed chain ankle DF ROM screen.

---

## CONFLICT OF INTEREST STATEMENT

Drs. Kiesel and Plisky have equity in Functional Movement Systems LLC who owns the rights to the ankle screen used in this study. Other authors have no conflicts of interest.

## ACKNOWLEDGEMENTS

Funding provided through the Ridgway Student Research Award from the University of Evansville

Submitted: July 13, 2020 CDT, Accepted: October 10, 2020 CDT



This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CCBY-NC-ND-4.0). View this license's legal deed at <https://creativecommons.org/licenses/by-nc-nd/4.0> and legal code at <https://creativecommons.org/licenses/by-nc-nd/4.0/legalcode> for more information.

## REFERENCES

1. Hootman JM, Dick R, Agel J. Epidemiology of collegiate injuries for 15 sports: Summary and recommendations for injury prevention initiatives. *J Athl Train*. 2007;42(2):311-319.
2. Dick R, Hootman JM, Agel J, et al. Descriptive epidemiology of collegiate women's field hockey injuries: National Collegiate Athletic Association Injury Surveillance System, 1988-1989 through 2002-2003. *J Athl Train*. 2007;42(2):211-220.
3. Lemoyne J, Poulin C, Richer N, et al. Analyzing injuries among university-level athletes: Prevalence, patterns and risk factors. *J Can Chiropr Assoc*. 2017;61(2):88-95.
4. Almeida SA, Williams KM, Shaffer RA, Brodine SK. Epidemiological patterns of musculoskeletal injuries and physical training. *Med Sci Sports Exerc*. 1999;31(8):1176-1182. doi:10.1097/00005768-199908000-00015
5. McKay GD, Goldie PA, Payne WR, et al. Ankle injuries in basketball: Injury rate and risk factors. *Br J Sports Med*. 2001;35(2):103-108. doi:10.1136/bjism.35.2.103
6. Nelson AJ, Collins CL, Yard EE, et al. Ankle injuries among United States high school sports athletes, 2005-2006. *J Athl Train*. 2007;42(3):381-387.
7. Cloke DJ, Spencer S, Hodson A, Deehan D. The epidemiology of ankle injuries occurring in English Football Association academies. *Br J Sports Med*. 2009;43(14):1119-1125. doi:10.1136/bjism.2008.052050
8. Malliaras P, Cook JL, Kent P. Reduced ankle dorsiflexion range may increase the risk of patellar tendon injury among volleyball players. *J Sci Med Sport*. 2006;9(4):304-309. doi:10.1016/j.jsams.2006.03.015
9. Pope R, Herbert R, Kirwan J. Effects of ankle dorsiflexion range and pre-exercise calf muscle stretching on injury risk in Army recruits. *Aust J Physiother*. 1998;44(3):165-172. doi:10.1016/s0004-9514(14)60376-7
10. Gabbe BJ, Finch CF, Wajswelner H, Bennell KL. Predictors of lower extremity injuries at the community level of Australian football. *Clin J Sport Med*. 2004;14(2):56-63. doi:10.1097/00042752-200403000-00002
11. Fong C-M, Blackburn JT, Norcross MF, McGrath M, Padua DA. Ankle-dorsiflexion range of motion and landing biomechanics. *J Athl Train*. 2011;46(1):5-10. doi:10.4085/1062-6050-46.1.5
12. Bastien M, Moffet H, Bouyer L, Perron M, Hébert LJ, Leblond J. Concurrent and discriminant validity of the Star Excursion Balance Test for military personnel with lateral ankle sprain. *J Sport Rehabil*. 2014;23(1):44-55. doi:10.1123/jsr.2012-0132
13. Martin RL, McPoil TG. Reliability of ankle goniometric measurements: A literature review. *J Am Podiatr Med Assoc*. 2005;95(6):564-572. doi:10.7547/0950564
14. Drewes LK, McKeon PO, Kerrigan DC, Hertel J. Dorsiflexion deficit during jogging with chronic ankle instability. *J Sci Med Sport*. 2009;12(6):685-687. doi:10.1016/j.jsams.2008.07.003
15. Hoch MC, Farwell KE, Gaven SL, Weinhandl JT. Weight-bearing dorsiflexion range of motion and landing biomechanics in individuals with chronic ankle instability. *J Athl Train*. 2015;50(8):833-839. doi:10.4085/1062-6050-50.5.07
16. Rabin A, Portnoy S, Kozol Z. The association of ankle dorsiflexion range of motion with hip and knee kinematics during the lateral step-down test. *J Orthop Sports Phys Ther*. 2016;46(11):1002-1009. doi:10.2519/jospt.2016.6621
17. Teyhen DS, Shaffer SW, Butler RJ, et al. What risk factors are associated with musculoskeletal injury in US Army Rangers? A prospective prognostic study. *Clin Orthop Relat Res*. 2015;473(9):2948-2958. doi:10.1007/s11999-015-4342-6
18. Bennell KL, Talbot RC, Wajswelner H, Techovanich W, Kelly D, Hall A. Intra-rater and inter-rater reliability of a weight-bearing lunge measure of ankle dorsiflexion. *Aust J Physiother*. 1998;44(3):175-180. doi:10.1016/s0004-9514(14)60377-9
19. O'Shea S, Grafton K. The intra and inter-rater reliability of a modified weight-bearing lunge measure of ankle dorsiflexion. *Man Ther*. 2013;18(3):264-268. doi:10.1016/j.math.2012.08.007
20. Tucker WS, Ingram RL. Reliability and validity of measuring scapular upward rotation using an electrical inclinometer. *J Electromyogr Kinesiol*. 2012;22(3):419-423. doi:10.1016/j.jelekin.2012.02.013

21. Gourlay J, Bullock G, Weaver A, et al. The reliability and criterion validity of a novel dorsiflexion range of motion screen. *Athl Train Sports Health Care*. 2020;12(1):40-44.
22. Munteanu SE, Strawhorn AB, Landorf KB, Bird AR, Murley GS. A weightbearing technique for the measurement of ankle joint dorsiflexion with the knee extended is reliable. *J Sci Med Sport*. 2009;12(1):54-59. [doi:10.1016/j.jsams.2007.06.009](https://doi.org/10.1016/j.jsams.2007.06.009)
23. Cicchetti DV. Guidelines, criteria, and rules of thumb for evaluating normed and standardized assessment instruments in psychology. *Psychol Assess*. 1994;6(4):284-290. [doi:10.1037/1040-3590.6.4.284](https://doi.org/10.1037/1040-3590.6.4.284)
24. Venturni C, André A, Aguilar BP, et al. Reliability of two evaluation methods of active range of motion in the ankle of healthy individuals. *Acta Fisiátrica*. 2006;13(1):39-43.
25. Krause DA, Cloud BA, Forster LA, Schrank JA, Hollman JH. Measurement of ankle dorsiflexion: A comparison of active and passive techniques in multiple positions. *J Sport Rehabil*. 2011;20(3):333-344. [doi:10.1123/jsr.20.3.333](https://doi.org/10.1123/jsr.20.3.333)
26. Denegar CR, Hertel J, Fonseca J. The effect of lateral ankle sprain on dorsiflexion range of motion, posterior talar glide, and joint laxity. *J Orthop Sports Phys Ther*. 2002;32(4):166-173. [doi:10.2519/jospt.2002.32.4.166](https://doi.org/10.2519/jospt.2002.32.4.166)
27. Driller MW, Overmayer RG. The effects of tissue flossing on ankle range of motion and jump performance. *Phys Ther Sport*. 2017;25:20-24. [doi:10.1016/j.ptsp.2016.12.004](https://doi.org/10.1016/j.ptsp.2016.12.004)
28. Dill KE, Begalle RL, Frank BS, Zinder SM, Padua DA. Altered knee and ankle kinematics during squatting in those with limited weight-bearing-lunge ankle-dorsiflexion range of motion. *J Athl Train*. 2014;49(6):723-732. [doi:10.4085/1062-6050-49.3.29](https://doi.org/10.4085/1062-6050-49.3.29)
29. Leanderson J, Wykman A, Eriksson E. Ankle sprain and postural sway in basketball players. *Knee Surg Sports Traumatol Arthrosc*. 1993;1(3-4):203-205. [doi:10.1007/bf01560207](https://doi.org/10.1007/bf01560207)