

Original Research

Utilizing Hip Abduction Strength to Body-Weight Ratios in Return to Sport Decision-Making After ACL Reconstruction

Steven Higbie, PT, DPT, SCS, CSCS^{1, a}, Jacquelyn Kleihege, PT, SCS¹, Brian Duncan, PT, DPT, SCS, OCS, FAAOMPT¹, Walter R. Lowe, MD², Lane Bailey, PhD, PT¹

¹ Sports Medicine and Rehabilitation, Memorial Hermann Ironman Sports Medicine Institute, ² Orthopaedic Surgery, McGovern Medical School at UT Health

Keywords: strength ratio, return to sport, hip abduction strength, acl injury

<https://doi.org/10.26603/001c.27346>

International Journal of Sports Physical Therapy

Vol. 16, Issue 5, 2021

Background

Despite the association between hip abduction weakness and non-contact anterior cruciate ligament (ACL) injury, hip abduction strength is rarely considered in return to sport decision-making following ACL reconstruction (ACLR).

Hypothesis/Purpose

The purpose of this study was to compare self-reported function, objective functional test performance, and re-injury rates in patients with high ($\geq 33\%$) versus low ($< 33\%$) isometric hip abduction strength to body weight (BW) ratios when returning to activity following ACLR.

Study Design

Cohort study

Methods

Data were gathered from a single-surgeon database and included baseline demographics. Clinical outcomes were assessed at the time of release to activity and included self-reported outcomes and a functional testing battery. Isometric hip abduction strength was obtained using a handheld dynamometer. Groups were dichotomized into those with low vs high strength to BW ratios. Two-year follow-up was performed using the single assessment numeric evaluation (SANE). Data were analyzed using univariate general linear models with an alpha level of .05.

Results

Of the 528 enrolled patients, 364 (68.9%) demonstrated a low strength to BW ratio. Baseline comparisons revealed more females and higher BMI ($P < .05$) in the $< 33\%$ group. At release to activity, the $< 33\%$ BW group demonstrated lower International Knee Documentation Committee survey scores (88.2 ± 13.6 vs 93.5 ± 10.3 , $P < .01$), ACL-Return to Sport After Injury (76.2 ± 15.4 vs 88.5 ± 16.9 , $P < .01$) scores, and isokinetic hamstring peak torque ($P = .04$). At 2-years, the $< 33\%$ group reported lower SANE scores (83.3 ± 21.1 vs 92.83 ± 11.4 , $P = .05$) with no significant differences in re-injuries.

Conclusion

Patients with low hip abduction strength to BW ratios demonstrated lower subjective function, psychological readiness, and isokinetic hamstring peak torque when completing functional testing following ACLR. Subjective deficits remained at 2-years.

Level of Evidence

Level 3

^a **Corresponding author:** Steven Higbie, PT, DPT, SCS, CSCS, Memorial Hermann Ironman Sports Medicine Institute, Department of Sports Medicine & Rehabilitation, Houston, TX, 77030, USA. Steven.higbie@memorialhermann.org

Key Terms

ACL injury, hip abduction strength, return to sport, strength ratio

Clinical Relevance

Assessing isometric hip abduction strength to body weight ratio may be beneficial in determining readiness to return to sport following ACL reconstruction.

What is Known About the Subject

Three prospective studies have provided conflicting evidence regarding the relationship between hip abduction strength and ACL injury. A clinical cut-point of hip abduction strength:BW ratio <35.4% has been suggested to identify athletes at risk of sustaining a non-contact ACL injury. To our knowledge no studies have examined isometric hip abduction strength:BW ratios in athletes attempting to return to sport following ACLR.

What This Study Adds to Existing Knowledge

This study examines the potential for hip abduction strength:BW ratio to be included as an additional metric in return to sport testing batteries.

INTRODUCTION

Non-contact injury mechanisms are responsible for approximately 70% of anterior cruciate ligament (ACL) ruptures.^{1,2} Several biomechanical and neuromuscular factors have been identified as risk factors for non-contact injuries,^{3,4} with prevention strategies often targeted at reducing dynamic knee valgus.⁵⁻⁷ Hip abductor strength may assist in controlling dynamic knee valgus by reducing hip adduction and internal rotation associated with dynamic knee valgus.^{4,8} Strengthening programs targeting the hip abductors are frequently included following ACL reconstruction (ACLR) to assist in normalizing lower-extremity strength and facilitating normal trunk, hip, and knee mechanics when returning to sport.^{9,10} However, despite the supporting evidence, few studies have included hip abduction strength as a functional criterion when determining readiness for return to sport.

Perhaps the omission of hip abduction strength from return-to-sport decision-making may be due to the inconsistencies found within the current literature and heterogeneity in testing procedures. Specifically, three prospective studies have provided conflicting evidence regarding the role of hip abduction strength in identifying athletes at risk for non-contact ACL injury. An investigation of 867 elite Norwegian female athletes demonstrated no association between hip abduction strength and ACL injury.¹¹ In contrast, an additional study suggested increased hip abduction strength was an independent risk factor for non-contact ACL injury in female Japanese high school basketball athletes.¹² The authors suggested that athletes with greater hip abduction strength may be predisposed to move into hip adduction in an attempt to counterbalance, highlighting the lack of consistent findings with regard to this clinical measure. Khayambashi et al.¹³ identified hip abduction weakness as an independent predictor of non-contact ACL injuries. In a study of 501 athletes, hip abduction weakness was significantly associated with ACL injury and explained 10.2% of the variation in injury status. A clinical threshold of hip abduction strength < 35.4% of body weight (BW) was reported to identify athletes at risk for future non-contact injury.¹³ The authors suggested using this hip ab-

duction strength to BW ratio (sensitivity 0.87, specificity 0.65) to assist in screening for ACL injury risk.¹³

Despite the potential link between hip abduction strength and ACL injury, little evidence exists regarding the inclusion of hip abduction strength in return to sport assessments. A recent systematic review of 209 studies identified only 86 studies utilizing strength measures as return to sport criteria, with minimal reporting of hip abduction strength.¹⁴ Several case-control and cross-sectional studies have compared hip abduction strength between healthy controls and ACLR patients who had already returned to sport,¹⁵⁻¹⁸ though none utilized hip abduction strength as a return to sport decision-making tool. To our knowledge, no studies have assessed the relationship between isometric hip abduction strength on a combined subjective and objective functional testing battery at time to return to sport. Further, no studies have examined if meeting a suggested hip abduction strength to body weight (BW) ratio is associated with improved outcomes following ACLR. Therefore, the purpose of this study was to compare self-reported function, objective functional test performance, and re-injury rates in patients with high ($\geq 33\%$) versus low ($< 33\%$) isometric hip abduction strength to BW ratios when returning to activity following ACLR. The authors hypothesized that individuals demonstrating hip abduction strength $\geq 33\%$ BW would perform better on self-reported outcomes, demonstrate superior objective functional testing performance, and demonstrate a lower re-injury rate when compared to those who exhibited lower strength ratios.

METHODS

A retrospective comparison study was conducted in accordance with the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) guidelines using a single surgeon (WRL) database of 829 patients who underwent ACLR from 2017-2019. All participants provided verbal and written consent. This study was approved by the University of Texas Health Sciences Center Institutional Review Board (HSC-MH-14-0734) and registered with [Clinicaltrials.gov](https://clinicaltrials.gov) (NCT03704376). The study inclusion criteria were patients who underwent ACLR, completed hip abduc-

tion strength testing and functional testing at time of release to activity, and completed follow-up surveys at two years post-surgery. Patients were excluded if they had orthopaedic conditions or medical complications preventing them from performing standard post-operative rehabilitation (fracture or deep vein thrombosis) or if they did not intend to return to sports participation. A total of 829 patients were reviewed in the database with 528 patients meeting inclusion criteria.

HIP STRENGTH-TO-BODYWEIGHT RATIOS

Bilateral isometric hip abduction strength testing was performed with the patient in supine and the hip in ten degrees of abduction (Figure 1). The testing leg was secured to the table with a belt placed proximal to the lateral femoral condyle. A HOGGAN microfet-2 handheld dynamometer (Hoggan Scientific, 3653 W. 1987 S., Salt Lake City, UT, USA) was secured between the belt and the testing leg. The patient was instructed to perform a maximum hip abduction contraction into the fixed resistance of the belt and examiner (make test) for 5 seconds. Standard verbal cueing instructions were used to minimize compensatory strategies. Three trials were completed and then averaged with the force being recorded in pounds. All testers completed reliability training and demonstrated excellent reliability with intraclass correlation coefficients ranging from 0.921-0.927 ($P < .01$). Patients were dichotomized into two groups for analysis: those demonstrating $<33\%$ hip abduction strength to BW ratio (low strength) and those $\geq 33\%$ (high strength). The cut-off of $\geq 33\%$ BW was used instead of the previously referenced 35.4% as it seemed more practical for clinical use. Specifically, $\geq 33\%$ BW may provide a notable threshold to aid in guiding clinical decision and counseling patient readiness to return to sport with the sports medicine team.

DATA COLLECTION

Baseline patient demographic characteristics were obtained for age, gender, height, weight, body mass index (BMI), and preinjury activity level via the Marx activity score.¹⁹ Subjective function at time of release to activity was assessed via the International Knee Disability Committee (IKDC-2000) survey²⁰ and the ACL Return to Sport Index (ACL-RSI)²¹ psychological readiness survey. Both groups were compared on functional test performance including passive knee range of motion (ROM), Y-balance anterior reach testing, four single-leg hop tests, and Biodex isokinetic quadriceps and hamstring testing. Y-balance (YBT) and hop test performance was assessed using previously established protocols^{22,23} with three trials being completed on the uninvolved and involved limbs. Isokinetic quadriceps and hamstring peak torque testing were completed via an isokinetic dynamometer (Biodex Medical Systems Inc., 20 Ramsey Rd., Shirley, NY, USA) at 60 (5 repetitions), 180 (10 repetitions), and 300 degrees (15 repetitions) per second ($^{\circ}/\text{sec}$). Patients completed serial functional testing until demonstrating $\geq 85\%$ quadriceps limb symmetry index (LSI) via isokinetic dynamometry, $\geq 90\%$ LSI on the remaining functional tests, and being cleared by the surgeon. Surgical data were reviewed from operative reports with compar-



Figure 1. Hip Abduction Strength Testing

Isometric hip abduction strength testing was completed in supine with the hip in 10 degrees of abduction. The patient was secured to the table via a stabilization belt. The patient completed a make test for 5 seconds with the peak force being recorded in pounds via a handheld dynamometer.

isons made based on graft type, procedure type, use of platelet-rich-plasma (PRP) or bone marrow aspiration (BMA), and type of rehabilitation protocol (accelerated or delayed). All patients followed a standardized accelerated or delayed rehabilitation protocol based on physician judgment at time of surgery. Patients in the delayed protocol group had protected weight-bearing and knee flexion ROM for the first four weeks following surgery. Two-year follow-up data was collected via electronic survey and included the Single Assessment Numeric Evaluation (SANE),²⁴ current level of sport participation, and re-injury status.

STATISTICAL ANALYSIS

Baseline patient demographic and surgical (graft type, use of BMA or PRP, rehabilitation protocol) information were analyzed via one-way (group) analysis of variance (ANOVA). Comparisons for subjective and objective functional testing scores at time of release to activity were analyzed via univariate analysis of variance. Between groups differences for all subjective reports at two-year follow-up were analyzed using an independent Student *t*-test. An *a priori alpha* of .05 was considered to be statistically significant for between groups' comparisons. All statistical analyses were performed using IBM SPSS Statistics (version 24, Armonk NY, USA) statistical software.

RESULTS

Of the 528 patients enrolled in the study, 364 (68.9%) had a ratio of $<33\%$ hip abduction strength to BW, and 164

Table 1. Patient Demographics

	Low Strength (n=364)	High Strength (n=164)	P Value
Age (years)	25.59 ± 12.36	22.84 ± 9.99	.031*
Gender (% male)	189 (51.9%)	103 (62.8%)	.036*
Height (inches)	68.03 ± 4.61	68.55 ± 4.10	.289
Weight (pounds)	172.61 ± 45.12	157.35 ± 30.91	.001*
BMI (kg/m ²)	26.03 ± 5.12	23.47 ± 3.45	.000*
MARX Score (0-16)	9.10 ± 5.48	10.39 ± 5.35	.347

Values reported as mean ± std dev. *Statistical Significance at $\leq .05$; Low Strength= patients demonstrating hip abduction strength: body weight (BW) ratio < 33%; High Strength= patients demonstrating hip abduction strength ratio $\geq 33\%$

Table 2. Subjective Scores and Functional Test Performance at Time of Release to Activity

	Low Strength (n=364)	High Strength (n=164)	P Value
IKDC (0-100)	88.15 ± 13.59	93.52 ± 10.27	.000*
ACL-RSI (0-100)	76.19 ± 15.39	88.52 ± 16.90	.000*
Extension ROM Deficit (deg.)	2.22 ± 1.75	2.38 ± 1.98	.432
Flexion ROM Deficit (deg.)	3.73 ± 4.41	3.86 ± 4.78	.782
Single Leg Balance Deficit (cm)	2.12 ± 4.13	2.19 ± 4.90	.892
Quadriceps LSI at 60 °/sec (%)	89.25 ± 20.24	86.55 ± 19.48	.149
Quadriceps LSI at 180 °/sec (%)	87.35 ± 17.60	87.88 ± 18.23	.511
Quadriceps LSI at 300 °/sec (%)	89.73 ± 15.94	89.77 ± 14.15	.723
Hamstring LSI at 60 °/sec (%)	94.38 ± 9.14	93.55 ± 6.27	.833
Hamstring LSI at 180 °/sec (%)	93.95 ± 7.03	96.23 ± 5.42	.035*
Hamstring LSI at 300 °/sec (%)	93.45 ± 7.37	99.93 ± 8.82	.041*
Single Leg Hop (LSI)	93.90 ± 6.86	94.04 ± 5.09	.248
Triple Hop (LSI)	92.37 ± 7.10	92.26 ± 7.37	.983
Cross-Over Hop (LSI)	91.29 ± 6.27	91.49 ± 6.95	.172
6m Timed Hop (LSI)	99.14 ± 1.14	99.13 ± 1.15	.681

Values reported as mean ± std dev. *Statistical Significance at $\leq .05$; Low Strength= patients demonstrating hip abduction strength: body weight (BW) ratio < 33%; High Strength= patients demonstrating hip abduction strength ratio $\geq 33\%$; IKDC= International Knee Documentation Committee Questionnaire; ACL-RSI= Anterior Cruciate Ligament Return to Sport After Injury Scale; ROM= Range of Motion; LSI= Limb Symmetry Index (involved limb/uninvolved limb).

(31.0%) exhibited a ratio $\geq 33\%$. [Table 1](#) outlines the baseline demographic characteristics of each cohort. Differences existed between the groups with the <33% BW group having a significantly lower percentage of males (51.9% vs 62.8%), older age (25.59 ± 12.36 vs 22.84 ± 9.99 years), higher weight (172.61 ± 45.12 vs 157.35 ± 30.91), and a higher BMI (26.03 ± 5.12 vs 23.47 ± 3.45). No significant differences existed for height or the Marx score ($P > .05$).

Subjective scores and functional test results at time to release to activity are shown in [Table 2](#). The <33% BW group reported lower IKDC (88.15 ± 13.59 vs 93.52 ± 10.3) and ACL-RSI (76.19 ± 15.39 vs 88.52 ± 16.90) scores. No significant differences were present between groups on ROM, Y-balance anterior reach testing, the single-leg hop tests, and three-speed quadriceps isokinetic testing ($P > .05$). Patients in the <33% BW group demonstrated significantly lower isokinetic hamstring peak torque LSI at 180°/sec (93.95 ± 7.03

vs 96.23 ± 5.42, $P = .035$) and 300°/sec (93.45 ± 7.37 vs 99.93 ± 8.82, $P = .041$). There were no significant differences in graft type, procedure type, surgical use of PRP or BMA, or rehabilitation protocol between groups. At two-year follow-up the <33% BW group reported significantly lower SANE scores (83.32 ± 21.06 vs 92.82 ± 11.36). No significant differences were observed for current level of sports participation ($P = .071$) and graft re-injury rate (4% vs 4%, $P = .986$) at two-years.

DISCUSSION

Hip abduction weakness is associated with altered jumping and landing mechanics⁸ and impaired running biomechanics.^{25,26} While these activities play a role in sports participation and returning to high-level activities, hip abduction

strength is rarely considered as a metric when considering an athlete's readiness to return to sport. Little evidence exists examining hip abduction strength following ACLR and its potential impact on subjective function and other objective measures of strength or stability. The current study is the first to our knowledge to demonstrate that athletes with low hip abduction strength to BW ratios report significantly lower subjective function and psychological readiness at time of release to activity and significantly lower subjective function at two-year follow-up.

A large percentage of patients in our study (68.9%) failed to demonstrate hip abduction strength $\geq 33\%$ BW. Significant demographic differences existed between groups with those demonstrating $<33\%$ BW being more female, older, heavier, and having a higher BMI. This aligns with previous work demonstrating a reduction in hip abductor isometric peak torque with aging²⁷ and gender differences.²⁸ Patients with higher BMI's may have struggled to achieve the cutoff strength ratio as they needed to reach larger raw strength values. Previous work has demonstrated that obese individuals demonstrate lower quadriceps peak torque:BW ratios when compared to leaner counterparts,²⁹ though little evidence has examined the impact of BMI on absolute and relative hip strength. Adjusting the 33% BW ratio based on gender or age may result in better identifying athletes with functional deficits at time of RTS.

In alignment with our hypothesis, those with low strength:BW ratios reported significantly lower psychological readiness and self-reported function at time of release to activity. Subjective deficits remained at two-year follow up with the $<33\%$ BW group reporting significantly lower function via the SANE score. Lower psychological readiness when returning to sport has been linked with second ACL injury³⁰ and failing to return to previous activity level.³¹ However, this study found no significant differences between re-injury rate and level of sport participation at two-year follow-up. Caution should be used when interpreting these results as the between group differences, though statistically significant, did not exceed the minimal clinical important difference for the IKDC-2000 or SANE score.³² Further, the ACL-RSI scores, though significantly lower in the $<33\%$ BW group, were above suggested cutoffs and previously reported average scores.^{33,34}

No significant between group differences were present for knee ROM, YBT performance, quadriceps isokinetic peak torque, and single leg hop testing in this study. This aligns with previous work concluding hip abduction strength was not predictive of single-leg hop performance following ACLR^{15,18} but differs from Clagg et al.³⁵ who found hip abduction strength was positively correlated with YBT reach distance. The subjective deficits observed in the $<33\%$ BW group may potentially be explained by hip abduction strength, as no significant differences existed between quad strength or functional performance. Patients in the $<33\%$ BW group had a significantly lower isokinetic hamstring

peak torque LSI at 180 and 300 °/sec. Hip abduction weakness may occur alongside posterior chain weakness and assist in explaining these deficits. Although significantly different, both groups demonstrated hamstring LSI values greater than the 90% value typically used for RTS clearance. Despite the $<33\%$ BW group reporting lower subjective function at two-year follow-up, no significant difference existed with re-injuries. This may be explained by both groups demonstrating $>85\%$ LSI for quad strength and $>90\%$ LSI for all hop testing.^{36,37}

Limitations of this study include the retrospective design and the lack of controlling for potential concomitant procedures (multi-ligament procedure, cartilage procedure, etc.) amongst the groups. The authors opted to use a cutoff score of 33% BW instead of the previously recommended 35.4% BW as it seemed for more practical for clinical practice. The use of a different BW ratio may serve as a more optimal cutoff to identify athletes who may perform more poorly at time of release to activity or two-year follow-up. Few previous studies exist for comparison with a wide heterogeneity in the methods used for hip abduction strength testing. The authors recommend future studies adopt a standardized, reliable methodology for assessing hip abduction strength. Future prospective studies would assist in determining the effect of hip abduction strengthening on outcomes following ACLR. Hip extensor strength,¹⁵ external rotation strength,^{13,18} and hip ROM³⁸ have been linked to ACL injury and should also be investigated in those returning to sport following ACLR.

CONCLUSIONS

Hip abduction strength to body weight ratio may provide helpful insight for clinical decision-making when determining when to release patients back to sport after ACL reconstruction. Patients failing to demonstrate hip abduction strength $\geq 33\%$ BW demonstrated lower psychological readiness, subjective function, and isokinetic hamstring peak torque at time of release to activity. Self-reported knee function remained significantly lower at two-year follow-up. Clinicians should consider the potential utility of hip abduction strength to body weight ratio when assessing readiness for return to sport following ACLR.

DISCLOSURES

Dr. Walter Lowe is a medical consultant of Don Joy Inc. and Arthrex Inc.

Submitted: March 09, 2021 CDT, Accepted: June 15, 2021 CDT



This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CCBY-NC-SA-4.0). View this license's legal deed at <https://creativecommons.org/licenses/by-nc-sa/4.0> and legal code at <https://creativecommons.org/licenses/by-nc-sa/4.0/legalcode> for more information.

REFERENCES

1. Boden BP, Dean GS, Feagin JA, Garrett WE. Mechanisms of anterior cruciate ligament injury. *Orthopedics*. 2000;23(6):573-578.
2. Krosshaug T, Nakamae A, Boden BP, et al. Mechanisms of anterior cruciate ligament injury in basketball: video analysis of 39 cases. *Am J Sports Med*. 2007;35(3):359-367. doi:10.1177/0363546506293899
3. Hewett TE, Di Stasi SL, Myer GD. Current concepts for injury prevention in athletes after anterior cruciate ligament reconstruction. *Am J Sports Med*. 2013;41(1):216-224. doi:10.1177/0363546512459638
4. Powers CM. The influence of abnormal hip mechanics on knee injury: a biomechanical perspective. *J Orthop Sports Phys Ther*. 2010;40(2):42-51. doi:10.2519/jospt.2010.3337
5. Padua DA, DiStefano LJ, Hewett TE, et al. National athletic trainers' association position statement: prevention of anterior cruciate ligament injury. *J Athl Train*. 2018;53(1):5-19. doi:10.4085/1062-6050-99-16
6. Hewett TE, Myer GD, Ford KR, Paterno MV, Quatman CE. Mechanisms, prediction, and prevention of ACL injuries: cut risk with three sharpened and validated tools. *J Orthop Res*. 2016;34(11):1843-1855. doi:10.1002/jor.23414
7. Nessler T, Denney L, Sampley J. ACL injury prevention: what does research tell us? *Curr Rev Musculoskelet Med*. 2017;10(3):281-288. doi:10.1007/s12178-017-9416-5
8. Paterno MV, Schmitt LC, Ford KR, et al. Biomechanical measures during landing and postural stability predict second anterior cruciate ligament injury after anterior cruciate ligament reconstruction and return to sport. *Am J Sports Med*. 2010;38(10):1968-1978. doi:10.1177/0363546510376053
9. Garrison JC, Bothwell J, Cohen K, Conway J. Effects of hip strengthening on early outcomes following anterior cruciate ligament reconstruction. *Int J Sports Phys Ther*. 2014;9(2):157-167.
10. Ferber R, Kendall KD, Farr L. Changes in knee biomechanics after a hip-abductor strengthening protocol for runners with patellofemoral pain syndrome. *J Athl Train*. 2011;46(2):142-149. doi:10.4085/1062-6050-46.2.142
11. Steffen K, Nilstad A, Kristianslund EK, Myklebust G, Bahr R, Krosshaug T. Association between lower extremity muscle strength and noncontact ACL injuries. *Med Sci Sports Exerc*. 2016;48(11):2082-2089. doi:10.1249/MSS.0000000000001014
12. Shimozaaki K, Nakase J, Takata Y, Shima Y, Kitaoka K, Tsuchiya H. Greater body mass index and hip abduction muscle strength predict noncontact anterior cruciate ligament injury in female Japanese high school basketball players. *Knee Surg Sports Traumatol Arthrosc*. 2018;26(10):3004-3011. doi:10.1007/s00167-018-4888-4
13. Khayambashi K, Ghoddosi N, Straub RK, Powers CM. Hip muscle strength predicts noncontact anterior cruciate ligament injury in male and female athletes: a prospective study. *Am J Sports Med*. 2016;44(2):355-361. doi:10.1177/0363546515616237
14. Burgi CR, Peters S, Ardern CL, et al. Which criteria are used to clear patients to return to sport after primary ACL reconstruction? A scoping review. *Br J Sports Med*. 2019;53(18):1154-1161. doi:10.1136/bjports-2018-099982
15. Tate J, Suckut T, Wages J, Lyles H, Perrin B. The associations between hip strength and hip kinematics during a single leg hop in recreational athletes post ACL reconstruction compared to healthy controls. *Int J Sports Phys Ther*. 2017;12(3):341-351.
16. Thomas AC, Villwock M, Wojtyls EM, Palmieri-Smith RM. Lower extremity muscle strength after anterior cruciate ligament injury and reconstruction. *J Athl Train*. 2013;48(5):610-620. doi:10.4085/1062-6050-48.3.23
17. Noehren B, Abraham A, Curry M, Johnson D, Ireland ML. Evaluation of proximal joint kinematics and muscle strength following ACL reconstruction surgery in female athletes: evaluations of kinematics and muscle strength. *J Orthop Res*. 2014;32(10):1305-1310. doi:10.1002/jor.22678
18. Kline PW, Burnham J, Yonz M, Johnson D, Ireland ML, Noehren B. Hip external rotation strength predicts hop performance after anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc*. April 2017. doi:10.1007/s00167-017-4534-6
19. Marx RG, Stump TJ, Jones EC, Wickiewicz TL, Warren RF. Development and evaluation of an activity rating scale for disorders of the knee. *Am J Sports Med*. 2001;29(2):213-218. doi:10.1177/03635465010290021601

20. Irrgang JJ, Anderson AF, Boland AL, et al. Development and validation of the international knee documentation committee subjective knee form. *Am J Sports Med.* 2001;29(5):600-613. doi:10.1177/03635465010290051301
21. Webster KE, Feller JA, Lambros C. Development and preliminary validation of a scale to measure the psychological impact of returning to sport following anterior cruciate ligament reconstruction surgery. *Phys Ther Sport Off J Assoc Chart Physiother Sports Med.* 2008;9(1):9-15. doi:10.1016/j.ptsp.2007.09.003
22. Garrison JC, Bothwell JM, Wolf G, Aryal S, Thigpen CA. Y balance TM test anterior reach symmetry at three months is related to single leg functional performance at time of return to sports following anterior cruciate ligament reconstruction. *Int J Sports Phys Ther.* 2015;10(5):602-611.
23. Ross MD, Langford B, Whelan PJ. Test-retest reliability of 4 single-leg horizontal hop tests. *J Strength Cond Res.* 2002;16(4):617-622.
24. Shelbourne KD, Barnes AF, Gray T. Correlation of a single assessment numeric evaluation (SANE) rating with modified Cincinnati knee rating system and IKDC subjective total scores for patients after ACL reconstruction or knee arthroscopy. *Am J Sports Med.* 2012;40(11):2487-2491. doi:10.1177/0363546512458576
25. Taylor - Haas JA, Hugentobler JA, DiCesare CA, et al. Reduced hip strength is associated with increased hip motion during running in youth adult and adolescent male long-distance runners. *Int J Sports Phys Ther.* 2014;9(4):456-467.
26. Luedke LE, Heiderscheid BC, Williams DSB, Rauh MJ. Association of isometric strength of hip and knee muscles with injury risk in high school cross country runners. *Int J Sports Phys Ther.* 2015;10(6):868-876.
27. Johnson ME, Mille M-L, Martinez KM, Crombie G, Rogers MW. Age-related changes in hip abductor and adductor joint torques. *Arch Phys Med Rehabil.* 2004;85(4):593-597. doi:10.1016/j.apmr.2003.07.022
28. Ramskov D, Pedersen MB, Kastrup K, et al. Normative values of eccentric hip abduction strength in novice runners: an equation adjusting for age and gender. *Int J Sports Phys Ther.* 2014;9(1):68-75.
29. Maffiuletti NA, Jubeau M, Munzinger U, et al. Differences in quadriceps muscle strength and fatigue between lean and obese subjects. *Eur J Appl Physiol.* 2007;101(1):51-59. doi:10.1007/s00421-007-0471-2
30. McPherson AL, Feller JA, Hewett TE, Webster KE. Psychological readiness to return to sport is associated with second anterior cruciate ligament injuries. *Am J Sports Med.* 2019;47(4):857-862. doi:10.1177/0363546518825258
31. Arden CL, Österberg A, Tagesson S, Gauffin H, Webster KE, Kvist J. The impact of psychological readiness to return to sport and recreational activities after anterior cruciate ligament reconstruction. *Br J Sports Med.* 2014;48(22):1613-1619. doi:10.1136/bjports-2014-093842
32. Winterstein AP, McGuine TA, Carr KE, Hetzel SJ. Comparison of IKDC and SANE outcome measures following knee injury in active female patients. *Sports Health.* 2013;5(6):523-529. doi:10.1177/1941738113499300
33. Sadeqi M, Klouche S, Bohu Y, Herman S, Lefevre N, Gerometta A. Progression of the psychological ACL-RSI score and return to sport after anterior cruciate ligament reconstruction: a prospective 2-year follow-up study from the French prospective anterior cruciate ligament reconstruction cohort study (FAST). *Orthop J Sports Med.* 2018;6(12). doi:10.1177/2325967118812819
34. Arden CL, Taylor NF, Feller JA, et al. Psychological responses matter in returning to preinjury level of sport after anterior cruciate ligament reconstruction surgery. *Am J Sports Med.* 2013;41(7):1549-1558. doi:10.1177/0363546513489284
35. Clagg S, Paterno MV, Hewett TE, Schmitt LC. Performance on the modified star excursion balance test at the time of return to sport following anterior cruciate ligament reconstruction. *J Orthop Sports Phys Ther.* 2015;45(6):444-452. doi:10.2519/jospt.2015.5040
36. Grindem H, Snyder-Mackler L, Moksnes H, Engebretsen L, Risberg MA. Simple decision rules can reduce reinjury risk by 84% after ACL reconstruction: the Delaware-Oslo ACL cohort study. *Br J Sports Med.* 2016;50(13):804-808. doi:10.1136/bjports-2016-096031
37. Toole AR, Ithurburn MP, Rauh MJ, Hewett TE, Paterno MV, Schmitt LC. Young athletes cleared for sports participation after anterior cruciate ligament reconstruction: how many actually meet recommended return-to-sport criterion cutoffs? *J Orthop Sports Phys Ther.* 2017;47(11):825-833. doi:10.2519/jospt.2017.7227
38. Boutris N, Byrne RA, Delgado DA, et al. Is there an association between noncontact anterior cruciate ligament injuries and decreased hip internal rotation or radiographic femoroacetabular impingement? A systematic review. *Arthrosc J Arthrosc Relat Surg.* 2018;34(3):943-950. doi:10.1016/j.arthro.2017.08.302